



EFFECT OF INTEGRATED NUTRIENT MANAGEMENT ON GROWTH AND YIELD OF INDIAN MUSTARD [*Brassica juncea* (L.) Czern. & coss.] UNDER IRRIGATED CONDITIONS OF PUNJAB

Ramanjot Kaur and Kamalesh Kumar

General Shivdev Singh Diwan Gurbachan Singh Khalsa College, Patiala, 147001

Email: jotzsaini30@gmail.com

Received: 22-05-2018

Accepted: 16-06-2018

ABSTRACT

Experiment was conducted during rabi season of 2015-16 at the Agricultural Research Farm Dhablan of the G.S.S.D.G.S. Khalsa college Patiala, Punjab. The experiment was laid out in randomized block design with 3 replications and 13 treatment combination of integrated nutrient Management. The soil of experimental field was clay in texture with pH 7.2 and contained organic carbon 0.80%, available nitrogen 374 kg/ha, available phosphorus 30.32 kg/ha and available K 120 kg/ha. All nutrients were applied in basal dose at one day before sowing. The crop was sown on 20th October, 2015. Application of Integrated nutrient Management significantly influenced the plant height, number of branches per plant, fresh weight of plant, dry weight of plant. The highest seed yield was recorded with the application of T7 (100% RDF + 2t FYM ha⁻¹ + 20kg S ha⁻¹ + 20kg ZnSO₄ ha⁻¹ + 1t vermicompost ha⁻¹ + Azotobacter Seed treatment) was found significantly higher over rest treatments.

Key word: Indian mustard, Integrated nutrient management, Siliquae.

Mustard is a major oilseed crop of India, it grown in tropical and temperate zones. They occupy a prominent place being next in importance to groundnut, both in area and production, meeting the fat requirement of about 50 percent population in India. It grown well from an altitude of 650 1500 meters, it require warm weather 20°C during seed germination, 15°C 20°C during plant growth and long sunny bright days 25°C 27°C at flowering and pod formation are most suitable for these crops. Mustard grow best under relatively cool temperatures upto flowering. These crops grow profusely at 30 60% relative humidity. Oilseeds have prestigious place in Indian agriculture next only to cereals. India is blessed with favourable agro-ecological conditions for the growth of wide range of cultivated, perennial and annual oilseeds.

Oilseeds are the most important crops in India both in respect of remunerative return per unit area and wider adaptability under constrained agro-climatic conditions. The oilseed crop Brassica are the third important edible oil sources after groundnut and soyabean, accounting for over 13.2 per cent of world's edible oil supply. In India, mustard and rapeseed are the second most important oilseed crops after groundnut contributing about 30 per cent of total oilseeds production Chand and Somani (2003).

Materials and Methods

A field experiment was carried out during Rabi season of 2015-16 at the Agricultural Research Farm Dhablan is situated at about 24-46 0N latitude and 76-24 0E longitude at an altitude of about 250 m above the mean sea level. The experiment was laid

out in randomized block design with 3 replications. From the five randomly selected plants the heights were recorded in cm. The numbers of branches were counted from the sample plants and the values of these were averaged. To study the fresh and dry weight of five plants were collected from the sampling rows of each plot at 30 days interval from sowing till harvest of the crop. These fresh samples were air dried and then dried in an oven at 60 °C till a constant weight was obtained and weighed to record the average dry weight of the plant. The weight of the sun dried harvested crop was recorded from net plot area and expressed in quintal per hectare after subtracting the seed yield. Seed yield of each plot excluding the border and sampling row was weighed in kilo gram and converted into quintal per hectare.

Result and Discussion

The use of integrated nutrient management is considered to be one of the most important factors to increase the plant height. The results of the present study showed that the application of integrated nutrient management statically significant by increase the plant height at various stages of recorded. The plant height increase with increasing fertilizer dose of INM treatment T7 (100% RDF + 2t FYM ha⁻¹ + 20kg S ha⁻¹ + 20kg ZnSO₄ ha⁻¹ + 1t vermicompost ha⁻¹ + Azotobacter Seed treatment) gives better response as comparison to other treatments. The beneficial effect of INM on plant height was also reported by Thaneshwar *et al.* (2017) in Indian mustard.

The result of the present study showed that the application of INM increase number of branches plant⁻¹ over control. The number of branch increased with increasing fertilizer dose levels of INM treatment seven showed better response as comparison to other treatments. This finding is also supported by Mohiuddin *et al.* (2011) Kumar *et al.* (2007), Singh and Kanaujia *et al.* (2009) in Indian

mustard. In general, INM addition increased the fresh and dry weight of the plant compared to control treatment. Then fresh and dry weight of plant increase with increasing fertilizer dose levels of INM treatment T7 (100% RDF + 2t FYM ha⁻¹ + 20kg S ha⁻¹ + 20kg ZnSO₄ ha⁻¹ + 1t vermicompost ha⁻¹ + Azotobacter Seed treatment) gives better response as comparison to other treatment.

Stover yield increased with increasing fertilizer dose levels of INM treatment seven gives better response as comparison to other treatments. The higher stover yield with the application of INM can be attributed to better growth of the plant as expressed in terms of plant height, number of branch plant⁻¹, fresh and dry weight of plant all above character responsible for stover yield. This finding is also supported by Singh and Kanaujia *et al.* (2009) and Mandal *et al.* (2006) in Indian mustard. The seed yield was affected by INM application. Such a positive yield response of INM application is obvious when it is limiting in the growing medium. Application of INM therefore provides better nutrition to Indian Mustard which resulted in higher seed yield. Increased in seed yield with the application of INM may be due to better growth of the plant as expressed in term of plant height, fresh. higher seed yield with the application of INM. This finding is also supported by Ramesh *et al.* (2009) in Indian mustard.

EFFECT OF INTEGRATED NUTRIENT MANAGEMENT ON GROWTH AND YIELD OF INDIAN MUSTARD
 [*Brassica juncea* (L.) Czern. & coss.] UNDER IRRIGATED CONDITIONS OF PUNJAB

Table:1 Effect of INM on plant height (cm) of Indian mustard at different stages of crop growth

Treatment	30 DAS	60 DAS	90 DAS	120DAS
T ₁	16.13	29.13	84.47	85.93
T ₂	19.27	35.13	87.87	90.20
T ₃	19.40	36.17	88.20	90.20
T ₄	20.53	38.00	89.57	91.57
T ₅	22.63	38.37	91.00	93.00
T ₆	23.60	45.10	91.87	93.87
T ₇	25.10	45.90	100.13	102.80
T ₈	19.20	32.90	86.63	88.63
T ₉	22.20	37.73	87.40	89.40
T ₁₀	22.47	36.93	88.70	90.70
T ₁₁	22.80	34.83	89.40	91.40
T ₁₂	23.03	37.53	89.27	91.27
T ₁₃	23.40	43.20	90.13	92.13
Mean	21.52	37.77	89.59	91.62
SE(d)±	NS	1.75	0.89	1.03
CD (5%)	5.94	3.96	2.00	2.33

Table: 2 Effect of INM on number of branches of Indian mustard at different stages of crop growth

Treatment	30 DAS	60 DAS	90 DAS	120 DAS
T ₁	1.13	2.33	3.73	5.07
T ₂	2.07	3.93	5.40	7.23
T ₃	2.47	4.13	5.73	7.73
T ₄	2.80	4.33	6.07	8.07
T ₅	3.33	4.53	6.53	8.53
T ₆	4.53	4.53	6.70	8.70
T ₇	5.67	6.93	6.93	9.00
T ₈	1.67	2.93	4.93	6.93
T ₉	2.60	3.87	5.87	7.87
T ₁₀	3.13	3.47	5.47	7.47
T ₁₁	3.13	3.80	5.60	7.60
T ₁₂	3.97	4.40	5.73	7.73
T ₁₃	4.33	5.27	5.87	7.80
Mean	3.14	4.19	5.74	7.67
SE(d)±	NS	1.28	0.88	0.69
CD (5%)	6.21	2.89	1.98	1.56

Table: 3 Effect of INM on number of leaves plant⁻¹ of Indian mustard at different stages of crop growth

Treatment	30 DAS	60 DAS	90 DAS	120DAS
T ₁	4.80	7.67	14.37	15.27
T ₂	6.53	8.60	17.37	19.37
T ₃	6.73	9.43	18.27	20.27
T ₄	7.33	10.93	19.73	21.73
T ₅	7.70	13.67	22.40	24.40
T ₆	9.00	14.17	22.77	24.77
T ₇	10.47	16.8	27.27	29.27
T ₈	7.07	7.87	16.53	18.53
T ₉	7.10	9.03	20.73	22.73
T ₁₀	7.27	9.93	21.07	23.07
T ₁₁	7.60	11.27	24.27	26.27
T ₁₂	8.53	12.77	25.87	27.87
T ₁₃	9.13	14.00	27.40	29.40
Mean	7.63	11.24	21.39	23.30
SE(d)±	0.49	1.11	1.86	1.69
CD (5%)	1.10	2.51	4.19	3.83

Table:4 Effect of INM on fresh weight (g plant⁻¹) of Indian mustard at different stages of crop growth

Treatment	30 DAS	60 DAS	90 DAS
T ₁	4.73	19.73	114.60
T ₂	6.13	21.60	117.37
T ₃	6.67	21.67	118.27
T ₄	6.07	23.03	119.73
T ₅	5.60	25.45	122.40
T ₆	6.93	33.26	122.77
T ₇	6.60	34.20	137.46
T ₈	4.60	20.18	116.53
T ₉	5.67	21.23	120.73
T ₁₀	5.80	22.93	121.07
T ₁₁	5.87	24.59	124.27
T ₁₂	6.43	24.79	125.87
T ₁₃	5.80	26.29	127.40
Mean	5.91	24.53	122.19
SE(d)±	0.80	1.78	1.50
CD (5%)	1.80	4.02	3.39

EFFECT OF INTEGRATED NUTRIENT MANAGEMENT ON GROWTH AND YIELD OF INDIAN MUSTARD
 [*Brassica juncea* (L.) Czern. & coss.] UNDER IRRIGATED CONDITIONS OF PUNJAB

Table: 5 Effect of INM on dry weight (g plant⁻¹) of Indian mustard at different stages of crop growth

Treatment	30 DAS	60 DAS	90 DAS
T ₁	1.40	8.33	29.73
T ₂	1.65	9.77	31.60
T ₃	2.17	9.53	31.67
T ₄	3.51	9.73	33.20
T ₅	2.57	9.87	35.45
T ₆	2.90	9.20	43.27
T ₇	4.30	9.60	45.20
T ₈	2.16	7.93	36.07
T ₉	2.75	8.67	31.23
T ₁₀	1.73	8.80	32.93
T ₁₁	2.49	8.80	34.59
T ₁₂	1.51	9.40	34.79
T ₁₃	2.93	8.87	36.29
Mean	2.47	9.11	35.08
SE(d)±	1.23	0.95	1.48
CD (5%)	2.79	2.14	3.34




Fig. 1: Effect of INM on dry weight (g) of Indian mustard

Table: 6 Effect of INM on stover and seed yield (q ha⁻¹) of Indian mustard

Treatment	Stover Yield	Seed Yield q ha ⁻¹
T ₁	11.37	9.07
T ₂	14.37	17.37
T ₃	15.27	19.27
T ₄	16.73	22.73
T ₅	19.40	24.07
T ₆	19.77	25.10
T ₇	24.27	26.93
T ₈	13.53	15.10
T ₉	17.73	17.40
T ₁₀	18.07	20.50
T ₁₁	21.27	22.60
T ₁₂	22.87	23.50
T ₁₃	24.40	25.05
Mean	18.39	20.66
SE(d)±	1.86	0.34
CD (5%)	4.19	0.76




Fig. No. 2: Effect of INM on stover and seed yield (q ha⁻¹) of Indian mustard

References

- Chand S and Somani LL (2003). Balanced use of fertilizers, organics and biofertilizers in mustard. *International Journal of Tropical Agriculture* 21 (1-4): 133-140.
- Mandal KG, Hati KM, Mishra AK and Bandyopadhyay KK (2006). Assessment of irrigation and nutrient effects on growth, yield and water use efficiency of Indian mustard (*Brassica juncea* L.) in central India. *Agricultural Water Management* 85 (3): 279-286.
- Mohiuddin M, Paul AK, Satradhar GNC, Bhuiyan MSI and Zubair HM (2011). Response of Nitrogen and sulphur fertilizer on yield, yield component and protein content of oilseed mustard (*Brassica spp.*) *International Journal of Bio-Resource and Stress Management*. 2(1): 93-99.
- Ramesh P, Panwar N R, Singh A B and Ramana S (2009). Effect of organic nutrient management practices on the production potential, nutrient uptake, soil quality, input-use efficiency and economics of mustard (*Brassica juncea* L.). *Indian Journal of Agricultural Sciences* 79 (1): 40-44.
- Kumar S, Sharma R and Dwivedi V K (2007). Response of Indian mustard (*Brassica juncea* L.) to organic based on distillery compost, farm yard manure and inorganic fertilizer. *Journal of Living World* 14 (2): 20-23.
- Singh P K, Imnuksungba and Kanajuia S P (2009). Effect of integrated nutrient management on growth, yield, its attributes and nutrients uptake of mustard crop in acidic soils of Nagaland. *Environment and Ecology* 27 (3): 1036-1039.
- Thaneshwar, Singh V, Jai Prakash, Kumar M, Kumar S and Singh RK (2017) Effect of Integrated Nutrient Management on Growth and Yield of Mustard (*Brassica juncea* L.) in Irrigated Condition of Upper Gangetic Plain Zone of India. *International Journal of Current Microbiology and Applied Sciences* 6 (1). 922-932.